



Macroplastique®

TREATMENT OF MALE
URINARY STRESS INCONTINENCE



Innovation for Health

PREDICTABLE PRODUCT. PROVEN PERFORMANCE.

MACROPLASTIQUE® BULKING AGENT

Macroplastique is indicated for stress urinary incontinence (SUI) often found in men following transurethral resection of the prostate (TURP) or radical prostatectomy.

- Demonstrated safety and efficacy
- Minimally invasive procedure
- Immediate results

MINIMALLY INVASIVE


- Significantly less invasive than the alternative treatment options
- Provides immediate relief of symptoms
- Fast recovery and minimal complications

PROVEN

- Clinical success since 1991
- Documented long-term success¹⁻⁷
 - Up to 5 year follow-up
 - Up to 80% success determined by urine loss (pad weight), reduction in pad usage, and assessment of patients' perception of the outcome of therapy

COST EFFECTIVE

Significantly less expensive than current alternative treatment methods.



1. Khastgir, J.N., & Shah, P.J.R. (2000). Macroplastique™ injections: How effective is it for stress incontinence in males? Proceedings ICS Annual Meeting, Tampere, Finland, August 28-31.
2. Arya, M., Hamid, R., Sharon, A., & Shah, P. (2002). The treatment of male stress incontinence in neuropathic bladders with polydimethylsiloxane. Proceedings ICS Annual Meeting, Heidelberg, Germany, August 28-30, 230.
3. Hamid, R., Arya, M., Khastgir, J., Patel, H., & Shah, P. (2003). The treatment of male stress urinary incontinence with polydimethylsiloxane in compliant bladders following spinal cord injury. *Spinal Cord*, 41, 286-289.
4. Pinggera, G.M., Furtmüller, B., Rehder, P., Fiala, H.R., Bartch, G., & Strasser, H. (2003). Submucosale blasenhalstunterspritzung mit Macroplastique® in der behandlung der stressinkontinenz nach radikaler prostatektomie. *Wiener Klin Wochenschrift. Middle Eur J Med*, 115(Suppl 1).
5. Herschorn, S., & Buencamino, D. (2002). Early results of intraurethral Macroplastique for post radical prostatectomy. *Neuro Urology*, 21(4), 332.
6. Mourad, S. (2003). Long term efficacy of Macroplastique injection for the treatment of urinary incontinence in males and females. *Europ Urol Supplements*, 2(1), 175.
7. Kymälä, T., Tainio, H., Raitanen, M., Teuvo, L.J., & Tammela, T.L. (2003). Treatment of postoperative male urinary incontinence using transurethral Macroplastique injections. *J Endourol*, 17(2), 113-115.

For complete instructions for use, storage, warnings, indications, contraindications, precautions, adverse reactions and disclaimer of warranties, please refer to the insert accompanying each Macroplastique. Models are for illustrative purpose only. Macroplastique is a registered trademark. Macroplastique is manufactured by Uroplasty LLC. ©2018 LABORIE. All rights reserved.

Distributed in Australia by
Endotherapeutics ABN 89 088 045 103
Level 1, 51 Rawson St, Epping NSW 2121
T: 1300 00 3636 F: +61 (2) 9869 7723
E: info@endotherapeutics.com.au
www.endotherapeutics.com.au



Endotherapeutics

EUROPE/INTERNATIONAL:
Tel.: +31 53 4803700
Email: info@laborie.com

HEADQUARTERS:
Tel.: +1 905 612 1170
Email: marketing@laborie.com



www.laborie.com